



Aloevera A Phytopharmacological Review

1. **Pratima indal rathod**, student vardhaman college of pharmacy
2. **Gaurav Manwar** assistant professor vardhaman college of pharmacy
3. **Nitin B kohale**, principal vardhaman college of pharmacy
4. **Suraj B Rathod** assistant professor, Vardhaman college of pharmacy

Date of Submission: 07-03-2023

Date of Acceptance: 19-03-2023

Abstract

Aloe vera has lengthly been used as a conventional medicinal drug for inducing wound healing. It is a herbal product that now a days is utilized in beauty industry. Benefits related to Aloe vera had been attributed to the polysaccharides contained withinside the gel of the leaves alevn though there are numerous symptoms for its use. Biological sports consist of promoting of wound healing, antifungal activity, anti-infl ammatory, anticancer and immunomodulatory. Gingival fi broblasts play an vital function in oral wound healing. Double blind-managed trials are wished to decide its actual effi cacy in oral health.

Key words: Aloe vera, oral health, polysaccharides, wound healing

I. Introduction

It is a stem much less or very short-stemmed plant developing to 80-a hundred cm tall, spreading via way of means of offsets and root sprouts. The leaves are thick and fleshy because of water garage tissue withi nside the leaves to live to tell the tale in dry regions of low rainfall. Leaves are inexperienced to grey- inexperienced, with a serrated margin. The flora are produced on a spike as much as ninety cm tall, every flower pendulous, with a yellow tubular corolla 2-three cm lengthly[1]. There had been numerous species below the genus Aloe, which includes Aloe vera, Aloe barbadensis, Aloe ferox, Aloe chinensis, Aloe indica, Aloe peyrii etc. Amongst these, Aloe vera Linn syn. Alo barbadensis Miller is commonplace unanimously as the accurate botanical supply of Aloe[2]. Aloe vera (Sanskrit-Ghrita kumari, Kumara;[3] Botanical-Aloe barbadensis Miller; Hindi-Guarpatha, Ghikanvar) is perennial succulent xerophyte, which develops water-garage tissue with inside the leaves to live to tell the tale in dry regions of low or erratic rainfall. The plant has stiff grey-inexperienced lance-shaped leaves containing clean gel in a valuable mucilaginous pulp. Benefits related with Aloevera

had been attributed to the polysaccharides contained with inside the gel of the leaves. Historically, it's been used for a number of medicinal purposes. Aloevera has lengthly been used as a conventional medicinal drug for inducing wound healing. Over the years, this plant has been regarded via way of means of some of names such as 'the wand of heaven', 'heaven's blessing' and 'the silent healer'. It is a herbal product that nowa days is used very a whole lot in beauty industry. Aloe Vera is derived from the tropical cactus of the genus aloe. Aloe Vera derives its call from the Arabic word Aloe which means shining sour substancel due to the sour liquid located with inside the leaves and Vera which means —truel in Latin [4, 5]. Aloevera is Aloe barbadensis miller. It belongs to Asphodelaceae (Liliaceae) family, and is a shrubby or arborescent, perennial, xerophytic, succulent, pea-inexperienced colour plant.The aloe plant has lengthly (up to twenty inches lengthly and five inches wide), triangular, fleshy leaves which have spikes alongside the edges. The sparkling parenchymal gel from the center of the leaf is clear; this element is from time to time dried to shape aloe Vera pay attention or diluted with water to create aloe juice products [6]. The sticky latex liquid is derived from the yellowish inexperienced pericyclic tubules that line the leaf (rind); this is the element that yields laxative anthraquinones [7]. There are over three hundred species of aloe, maximum of which might be local to South Africa, Madagascar and Arabia [8]. The distinctive species have fairly distinctive concentrations of lively ingredients Aloe Vera is one of the oldest medicinal flora for wholesome pores and skin ever known. This plant is regularly stated utilized in natural drug treatments due to the fact the start of the first century AD [10]. Aloe Vera (AV) gel has been used withinside the remedy of wounds, burns, insect stings, and pores and skin Inflammation, anti-inflammatory, antiseptic and antimicrobial, anti-tumor, anti-pores and skin protection, anti-diabetic, anti-bacterial, anti-viral, and whicllare verycritical for wound recovery [11, 12]. AV is powerful for wound recovery through diverse mechanisms which



include keeping wet wounds, growing mobileular migration, growing collagen production, and decreasing inflammation [13].

Active Ingredient of Aloe vera

More than 75 active ingredients from inner gel have been identified including vitamins, minerals, enzymes, sugars, anthraquinones or phenolic compounds, lignin, saponins, sterols, amino acids and salicylic acid. Active ingredients of Aloe vera leaf pulp and exudates[14].

Vitamins :-carotene), choline, folic acid, -tocopherol

Enzymes :- Alkaline phosphatase, amylase, carboxypeptidase, catalase, bradikininase, cyclooxygenase, peroxidase, carboxy-peptidase, cyclooxygenase, lipase, oxidase, phosphoenolpyruvate carboxylase, superoxide dismutase Anthraquinones/ anthrones Aloe-emodin, aloetic-acid, anthranol, aloin A and B (or collectively known as barbaloin), isobarbaloin, emodin, ester of cinnamic acid

Inorganic compounds:- Calcium, chlorine, chromium, copper, iron, magnesium, manganese, selenium, zinc, potassium, phosphorous, sodium

Carbohydrates :- Pure mannan, acetylated mannan, acetylated glucomannan

(acemannan), galactan, glucogalactomannan, galactogalacturan, galactoglucoarabinomannan, arabinogalactan, pectic substance, xylan, cellulose Saccharides:- Mannose, glucose, L-rhamnose, aldopentose

Organic compounds and lipids :- -linolenic acid, steroids (campesterol, cholesterol, -sitosterol), triglycerides, triterpenoid, gibberillin, lignins, potassium sorbate, salicylic acid, uric acid Chromones 8-C-glucosyl-

O-cinnamoyl)-7-O-methylaloeidin A, 8-C-glucosyl-(S)- aloesol, 8-C-glucosyl-7-O-methyl-(S)-aloesol, 8-C-glucosyl-7-O-methylaloeidin, 8-C-glucosyl-noreugenin, isoaloeserin D,

Isorabaichromone Nonessential and essential amino acids :- Alanine, arginine, aspartic acid, glutamic acid, glycine, histidine, hydroxyproline, isoleucine, leucine, lysine, methionine, phenylalanine, proline, threonine, tyrosine, valine.

Phyto pharmacological Action

Anti-inflammatory effects

It inhibits the cyclooxygenase pathway and decreases prostaglandin E2. Recently, the radical anti inflammatory compound called C-glycosyl chromone became removed from gel extracts.[15] Recently, the peptidase bradykinase became removed from aloe and proven to interrupt down the brady

kinin, an inflammatory substance that induces pain.[16]

Antibacterial

The interest of Aloe vera internal gel in opposition to each Gram-positive And Gram-poor micro organism has been verified with the aid of using several One-of-a-kind methods.[17] Streptococcus pyogenes and Streptococcus Faecalis are microorganisms which have been inhibited by with the aid of using Aloe vera gel.[18] Aloe vera gel reportedly changed into in opposition to Pseudomonas aeruginosa at the same time as acemannan avoided it from adhering to human lung epithelial cells in a monolayer culture.[19]

Antifungal

A processed Aloe vera gel preparation reportedly inhibited the growth of Candida albicans.[18]

Antiviral effects

This motion can be direct and indirect. Indirect due to stimulation of immune machine and direct is due to anthraquinones. The anthraquinones aloin turns on various enveloped virus; herpes simplex, varicella zoster and influenza.[20] Effect on immune machine Immunomodulating outcomes arise through activation of macrophage cells to generate nitric oxide, secrete cytokines (e.g., tumor necrosis factor-1, interleukin-6 and interferon-and gift mobileular floor markers.[21,22]

Antioxidant property

Glutathione peroxides activity, superoxide dismutase enzymes and a phenolic antioxidant have been located to be found in Aloe vera gel, which can be liable for those antioxidant effects.[23]

Antitumor effect

The fractions from aloes which might be claimed to have anticancer consequences encompass glycoproteins (lectins) and polysaccharides. Different research indicated antitumor interest for Aloe vera gel in phrases of decreased tumor burden, tumor shrinkage, tumor

necrosis and extended survival rates. An induction of glutathione S-transferase and an inhibition of the tumor-selling impact of phorbol myristic acetate has additionally been mentioned which recommend aloe gel in cancer chemoprevention.[24] Indirect motion on antitumor activity is stimulation of the immune response.[25]

Laxative effect (Purgative effect)

Anthraquinones increase intestine water content, stimulate water secretion and increase intestinal peristalsis.[26]

Antiseptic Properties



Aloe vera contain six antiseptic agent; lupeol, salicylic acid, urea Nitrogen, cinnamonic acid, phenol and sulfur.[27]

Effect On Gastric Acid Secretion And Ulcer

Aloevera gel has the capacity to therapy gastric ulcers or defend towards its formation in each animals and humans. The anti-ulcer sports of A. vera have been attributed to numerous feasible mechanisms consisting of its anti-inflammatory properties, restoration effects, mucus stimulatory effects, law of gastric secretions and presence of lectins[28]. The lectins inhibit aminopyrine uptake with the aid of using parietal cells. Thus, this ordinary capacity of the extract to inhibit gastric acid output may be a end result of direct movement at the acid generating cells[29].

II. Conclusion

Aloevera gel has the cappotential to treatment gastric ulcers or defend in opposition to its formation in each animals and humans. The anti-ulcer sports of A. vera have been attributed to numerous feasible mechanisms which include its anti-inflammatory properties, restoration effects, mucus stimulatory effects, law of gastric secretions and presence of lectins[28]. The lectins inhibit aminopyrine uptake with the aid of using parietal cells. Thus, this ordinary cappotential of the extract to inhibit gastric acid output can be a end result of direct movement at the acid generating cells[29].

Reference

- [1]. Kumar KPS, Bhowmik D, Chiranjib and Biswajit, Aloe vera: A Potential Herb And its Medicinal Importance, Journal of Chemistry and Pharmaceutical Research, 2(1), 2010, 21-29.
- [2]. Chandegara VK, Varshney AK, Aloe vera L. Processing and products: A Review. International Journal of Medicinal and Aromatic Plants 3(4), 2013, 492-506.
- [3]. Ayurveda-Recipes.com/Aloe_vera. [Homepage on the Internet]. Accessed on 2011 July 4. Available from: <http://www.ayurveda-recipes.com/aloevera.html>
- [4]. Gage, D. (1996). Aloe vera: Natures Soothing Healer. Healing Acts Press, Rochester, Vermont, USA, 120 p. 3.
- [5]. Surjusha, A., Vasani, R., & Saple, D. G. (2008). Aloe vera: a short review. Indian journal of dermatology, 53(4), 163-166.
- [6]. Schulz, V., Hansel, R., & Tyler, V. E. (1997). Rational Phyto therapy: A Physicians' Guide to Herbal Medicine. Berlin: Springer, 306.
- [7]. Christaki, E. V., & Florou-Paneri, P. C. (2010). Aloe vera: a plant for many Uses. J Food Agric Environ, 8(2), 245-249.
- [8]. Yagi, A., Tsunoda, M., Egusa, T., Akasaki, K., & Tsuji, H. (1998). Immunochemical distinction of Aloe vera, A. arborescens, and A. chinensis Gels. Planta medica, 64(03), 277-278.
- [9]. Van Wyk, B. E., Van Oudtshoorn, M. V. R., & Smith, G. F. (1995). Geographical variation in the major compounds of Aloe ferox leaf exudate. Planta Medica, 61(03), 250-253.
- [10]. Saito, M., Tanaka, M., Misawa, E., Yao, R., Nabeshima, K., Yamauchi, K., & Furukawa, F. (2016). Oral administration of Aloe vera gel powder prevents UVB-induced decrease in skin elasticity via suppression of overexpression of MMPs in hairless mice. Bioscience, biotechnology, and biochemistry, 80(7), 1416-1424.
- [11]. García, M. H., Juárez, J. A. T., & Jiménez, A. D. (2019). Importance and Properties of Aloe vera in the Production of Hair Shampoo. The Journal of Middle East and North Africa Sciences, 5(7), 18-23.
- [12]. Guenther, L., Lynde, C. W., Andriessen, A., Barankin, B., Goldstein, E., Skotnicki, S. P., & Sloan, K. (2012). Pathway to dry skin prevention and Treatment. Journal of cutaneous medicine and surgery, 16(1), 23-31.
- [13]. Komatsu, D., Mistura, D. V., Motta, A., Domingues, J. A., Hausen, M. A., & Duek, E. (2017). Development of a membrane of poly (L-co-D, L lactic acid-Co-trimethylene carbonate) with aloe vera: An alternative biomaterial designed To improve skin healing. Journal of biomaterials applications, 32(3), 311-320.
- [14]. Josias H. Hamman composition and applications of aloe vera leaf gel. Molecules 2008;13:1599-616.
- [15]. Hutter JA, Salmon M, Stavinoha WB, Satsangi N, Williams RF, Streeper RT, et al. Anti-inflammatory C-glycosyl chromone from aloe barbadensis. J Nat Prod 1996;59:541-3.
- [16]. Ito S, Teradaira R, Beppu H, Obata M, Nagatsu T, Fujita K. Properties and Pharmacological activity of carboxypeptidase in Aloe arborescens Mill var. Natalensis Berger. Phytother Res 1993;7:S26-9.



- [17]. Habeeb F, Shakir E, Bradbury F, Cameron P, Taravati MR, Drummond AJ, et Al. Screening methods used to determine the anti-microbial properties of Aloe Vera inner gel. *Methods* 2007;42:315-20.
- [18]. Heggors JP, Pineless GR, Robson MC. Dermaide aloe/aloe vera gel: Comparison of the antimicrobial effects. *J Am Med Technol* 1979;41:293-4.
- [19]. Azghani AO, Williams I, Holiday DB, Johnson AR. A beta-linked mannan Inhibits adherence of *Pseudomonas aeruginosa* to human lung epithelial cells. *Glycobiology* 1995;5:39-44.
- [20]. Sydiskis RJ, Owen DG, Lohr J, Rosler KH, Blomster RN. Inactivation of Enveloped viruses by anthraquinones extracted from plant. *Antimicrob Agent Chemother* 1991;35:2463-6.
- [21]. Tai-Nin Chow J, Williamson DA, Yates KM, Goux WJ. Chemical Characterisation of the immunomodulating polysaccharide of Aloe vera L. *Carbohydr Res* 2005;340:1131-42.
- [22]. Im SA, Oh ST, Song S, Kim MR, Kim DS, Woo SS, et al. Identification of Optimal molecular size of modified Aloe polysaccharides with maximum Immunomodulatory activity. *Int Immunopharmacol* 2005;5:271-9.
- [23]. Khan MA, Tania M, Zhang D, Chen H. Antioxidant enzyme and cancer. *Chin J Cancer Res* 2010;22:87-92.
- [24]. Kim HS, Lee BM. Inhibition of benzo (a) pyrene-DNA adduct formation by Aloe barbedebsis miller. *Carcinogenesis* 1997;18:771-6.
- [25]. Steenkamp V, Stewart MJ. Medicinal applications and toxicological activities Of Aloe products. *Pharm Biol* 2007;45:411-20.
- [26]. Ishii Y, Tanizawa H, Takino Y. Mechanism of cathartic effect. *Biol Pharm Bull* 1994;17:651-3.
- [27]. Surjushe A, Vasani R, Saple DG. Aloe vera: A short review. *Indian J Dermatol* 2008;53:163-6.
- [28]. Hamman JH, Composition and Applications of Aloe vera Leaf Gel, *Molecules*, 13, 2008, 1599-1616.
- [29]. Bhuvana KB, Hema NG, Patil RT, Review on aloe vera, *International Journal Of Advanced Research*, 2(3), 2014, 677-691.